

## Vibrational Spectrum and Normal Coordinate Analysis of *p*-Toluidino-*p*-chlorophenylglyoxime

HAKAN ARSLAN\* and TALAT ÖZPOZAN†

*Department of Chemistry, Faculty of Arts and Science*

*Mersin University, Mersin 33343, Turkey*

*Tel: (90)(532)7073122; Fax: (90)(324)3415759*

*E-mail: arslanh@mersin.edu.tr*

A complete vibrational assignment of the solid-state IR spectrum of *p*-toluidino-*p*-chlorophenylglyoxime was performed on the basis of normal coordinate analysis of a single molecule, assuming  $C_s$  point group symmetry. The calculated normal frequencies were in good agreement with the experimental one.

**Key Words:** *Vic*-dioxime, FTIR spectra, Valence force fields calculation, Normal coordinate analysis.

### INTRODUCTION

*Vic*-dioximes have an important place among coordination compounds. The early studies on oxime complexes started in 1905 as mentioned in review of Chakravorty<sup>1</sup> where the reactions of dimethylglyoximes with Ni(II) ion were examined. Later in 1907, Tschugaff<sup>2</sup> isolated dimethylglyoxime complexes of Co(III) which played an important role as a model compound in the elucidation of some biological and biochemical mechanisms. Several studies were concentrated on *vic*-dioximes complexed with Co atom in the following years<sup>3,4</sup>. They were used as model compounds to explain the structure of vitamin B12 and coenzyme B12 since these big molecules play important roles in the biological systems.

At present, new interesting properties of *vic*-dioximes are under examination. In some studies, its Pt complexes were used as anti-tumor agents in chemotherapy of some certain types of cancer. Some were used in the production of semi-conductors. In some other studies, liquid crystal properties of some *vic*-dioximes were being examined<sup>5</sup>. They were also used as column packing materials in chromatographic separation of nucleotides and nucleosides after bonding to natural resins as functional groups<sup>6</sup>.

Vibrational spectra of dioximes were not examined before but the some metal complexes of the glyoxime molecule were studied by using only its IR spectrum<sup>7</sup>. Since these classes of compounds have a tendency of fluorescing, it is difficult to take their Raman spectra and use it in normal coordinate analysis (NCA).

Based upon the literature search, a complete vibrational analysis of this type of *vic*-dioximes is not established so far. Having such important properties of

†Department of Chemistry, Faculty of Arts and Science, Erciyes University, Kayseri, Turkey.

*vic*-dioxime derivatives, *p*-toluidino-*p*-chlorophenylglyoxime (pTpCPG), a complete vibrational analysis using FTIR measurement data along with the result of normal coordinate analysis are presented in this paper.

## EXPERIMENTAL

*p*-Toluidino-*p*-chlorophenylglyoxime (pTpCPG) was synthesized as previously described in the literature<sup>8</sup>. Spectroscopically pure chemicals were obtained from Aldrich Chemicals, U.S.A. and used as such for recording spectra. The solid-state infrared spectrum of the pTpCPG studied were recorded using in the form of KBr pellet by BOMEN MB102 FTIR instrument in 4000–200  $\text{cm}^{-1}$  frequency ranges; the resolution was 1  $\text{cm}^{-1}$ . The Raman spectrum of the compound in a spinning cell was excited using the 488.0 nm line of  $\text{Ar}^+$  gas laser and recorded on a Jobin-Yvon U 1000 spectrometer, which was calibrated against the laser plasma emission lines.

### Normal coordinate analysis

Fig. 1 shows the structure of a *p*-toluidino-*p*-chlorophenylglyoxime molecule with the labeling of atoms. No structural studies are available for pTpCPG in the literature. Therefore, the structural parameters have been taken from related small molecules<sup>7, 9–15</sup>. The molecular parameters used for the calculations are reported in Table-1. Owing to its structure, the molecule pTpCPG belongs to the  $C_s$  symmetry point group. The 99 normal modes of pTpCPG are distributed between the two species ( $A'$  and  $A''$ ) of the  $C_s$  point group as:  $66A' + 33A''$ . All of the species are both IR and Raman active.




Fig. 1. Structure of *p*-toluidino-*p*-chlorophenylglyoxime

TABLE-1  
MOLECULAR PARAMETERS USED FOR THE CALCULATIONS

Bond Lengths (Å)			
C <sub>1</sub> —C <sub>2</sub>	1.435	N <sub>3</sub> —H <sub>14</sub>	1.008
C <sub>2</sub> —N <sub>3</sub>	1.472	O <sub>13</sub> —H <sub>15</sub>	0.956
C <sub>1</sub> —N <sub>4</sub>	1.290	C <sub>7</sub> —H <sub>17</sub>	1.084
C <sub>1</sub> —C <sub>6</sub>	1.510	C <sub>9</sub> —Cl <sub>19</sub>	1.700
C <sub>6</sub> —C <sub>7</sub>	1.397	N <sub>3</sub> —C <sub>22</sub>	1.426
N <sub>4</sub> —O <sub>12</sub>	1.346	C <sub>25</sub> —C <sub>32</sub>	1.510
Bond Angles (°)			
C <sub>2</sub> —C <sub>1</sub> —N <sub>4</sub>	125.42	C <sub>6</sub> —C <sub>7</sub> —H <sub>17</sub>	120.00
C <sub>2</sub> —C <sub>1</sub> —C <sub>6</sub>	117.79	C <sub>8</sub> —C <sub>9</sub> —Cl <sub>19</sub>	120.00
C <sub>2</sub> —N <sub>3</sub> —H <sub>14</sub>	120.55	N <sub>4</sub> —O <sub>12</sub> —H <sub>16</sub>	97.95
C <sub>2</sub> —N <sub>3</sub> —C <sub>22</sub>	118.90	N <sub>3</sub> —C <sub>22</sub> —C <sub>27</sub>	120.00
C <sub>1</sub> —N <sub>4</sub> —O <sub>12</sub>	121.45	C <sub>25</sub> —C <sub>32</sub> —H <sub>33</sub>	109.47
C <sub>1</sub> —C <sub>6</sub> —C <sub>7</sub>	120.00	H <sub>34</sub> —C <sub>32</sub> —H <sub>35</sub>	109.47

In order to ascertain the amount of mixing among the internal coordinates and to obtain a more accurate description of the fundamental vibrations of pTpCPG, a normal coordinate calculation has been undertaken. The calculations carried out using Simple General Valence Force Field (SGVFF) in Wilsons GF matrix method with the computer program originally written by Schachtschneider<sup>16</sup> and developed (for OS/2 -IBM) by Fischer *et al.*<sup>17</sup> under the name of SPSIM (SPectrum SIMultation).

The initial set of valence force constant and the corresponding off diagonal constant were transferred from related systems<sup>11-13, 18-28</sup>. A zero order calculation with the transferred force constant was performed except for some deformational and longitudinal modes. The results indicated the reasonable agreement between the calculated and the observed frequencies. The initial set of force constant was refined by the method of least square technique by keeping some interaction force constants fixed throughout the refinement process. The final values of force constants and their description are given in Table-2. The calculated and observed wavenumber values are compared in Table-3, which also give the significant values of the potential energy distribution (PED) for each mode of vibration.

## RESULTS AND DISCUSSION

The vibrational analysis of the pTpCPG under examination was performed in the foregoing study. FTIR part of the vibrational spectra of the compound were the only spectral source because of the difficulty of taking dispersive Raman spectra of the compound excited with a visible Laser source. A broad photoluminescence band along the IR wavelength region from 200 to 3500 cm<sup>-1</sup> prevented obtaining a reasonable Raman signals. This was not a serious problem in the compound because of both A' and A'' modes are active in both Raman and infrared in C<sub>s</sub> point group symmetry. The IR spectrum of the *p*-toluidino-*p*-chlorophenylglyoxime molecule is given in Fig. 2.


Fig. 2. Infrared spectrum of *p*-toluidino-*p*-chlorophenylglyoxime

TABLE-2  
 VALENCE FORCE CONSTANTS OF *p*-TOLUIDINO-*p*-  
 CHLOROPHENYLGLYOXIME†‡§

No	Force constant		Internal coordinate
1	$K(C_{ox}-C_{ox})$	5.240	$C_1-C_2$
2	$K(C=N)$	9.549*	$C_1-N_4$
3	$K(C_{ox}-C_x)$	5.101	$C_1-C_6$
4	$K(C_{ox}-N)$	6.116	$C_2-N_3$
5	$K(N-H)$	6.247*	$N_3-H_{14}$
6	$K(N-C_l)$	4.500	$N_3-C_{22}$
7	$K(N-O)$	4.010	$N_4-O_{12}$
8	$K(C-C)_{x, o}$	6.415	$C_6-C_7$
9	$K(C-C)_{x, m}$	6.727	$C_7-C_8$
10	$K(C-H)_x$	5.076*	$C_7-H_{17}$
11	$K(C-C)_{x, p}$	6.760	$C_8-C_9$
12	$K(C-Cl)$	3.733	$C_9-Cl_{19}$
13	$K(O-H)$	6.103*	$O_{12}-H_{16}$
14	$K(C-C)_l$	6.415	$C_{22}-C_{23}$
15	$K(C-H)_l$	4.999*	$C_{23}-H_{28}$
16	$K(C-CH_3)$	4.681	$C_{25}-C_{32}$
17	$K(C_l-H)$	4.588*	$C_{32}-H_{33}$
18	$H(C_{ox}-C_{ox}=N)$	2.651	$C_2-C_1-N_4$
19	$H(C_{ox}-C_{ox}-C_x)$	0.350	$C_2-C_1-C_6$
20	$H(N-C_{ox}-C_x)$	2.651	$N_4-C_1-C_6$

No	Force constant		Internal coordinate
21	H(C <sub>ox</sub> —C <sub>ox</sub> —N)	0.350	C <sub>1</sub> —C <sub>2</sub> —N <sub>3</sub>
22	H(C <sub>ox</sub> —C <sub>ox</sub> =N)	2.651	C <sub>1</sub> —C <sub>2</sub> —N <sub>5</sub>
23	H(N—C <sub>ox</sub> =N)	1.654*	N <sub>3</sub> —C <sub>2</sub> —N <sub>5</sub>
24	H(C <sub>ox</sub> —N—H)	0.901	C <sub>2</sub> —N <sub>3</sub> —H <sub>14</sub>
25	H(C <sub>ox</sub> —N—C <sub>1</sub> )	1.177	C <sub>2</sub> —N <sub>3</sub> —C <sub>22</sub>
26	H(H—N—C <sub>1</sub> )	0.901	H <sub>14</sub> —N <sub>3</sub> —C <sub>22</sub>
27	H(C <sub>ox</sub> —C—C) <sub>x</sub>	0.716	C <sub>1</sub> —C <sub>6</sub> —C <sub>7</sub>
28	H(C—C—C) <sub>x</sub>	1.242	C <sub>7</sub> —C <sub>6</sub> —C <sub>11</sub>
29	H(C—C—C) <sub>x, o</sub>	0.989	C <sub>6</sub> —C <sub>7</sub> —C <sub>8</sub>
30	H(C—C—H) <sub>x</sub>	0.508	C <sub>6</sub> —C <sub>7</sub> —H <sub>17</sub>
31	H(C—C—C) <sub>x, m</sub>	0.865	C <sub>8</sub> —C <sub>9</sub> —C <sub>10</sub>
32	H(C—C—Cl)	0.828	C <sub>8</sub> —C <sub>9</sub> —Cl <sub>19</sub>
33	H(C <sub>ox</sub> =N—O)	1.180	C <sub>1</sub> —N <sub>4</sub> —O <sub>12</sub>
34	H(N—O—H)	0.682*	N <sub>4</sub> —O <sub>12</sub> —H <sub>16</sub>
35	H(N—C <sub>1</sub> —C <sub>1</sub> )	0.760	N <sub>3</sub> —C <sub>22</sub> —C <sub>23</sub>
36	H(C—C—C) <sub>1</sub>	1.248	C <sub>23</sub> —C <sub>22</sub> —C <sub>27</sub>
37	H(C—C—C) <sub>1, o</sub>	1.028	C <sub>22</sub> —C <sub>23</sub> —C <sub>24</sub>
38	H(C—C—H) <sub>1</sub>	0.515	C <sub>22</sub> —C <sub>23</sub> —H <sub>28</sub>
39	H(C—C—CH <sub>3</sub> )	0.754	C <sub>24</sub> —C <sub>25</sub> —C <sub>32</sub>
40	H(C—C <sub>1</sub> —H <sub>1</sub> )	0.642	C <sub>25</sub> —C <sub>32</sub> —H <sub>33</sub>
41	H(H—C <sub>1</sub> —H)	0.535	H <sub>33</sub> —C <sub>32</sub> —H <sub>34</sub>
42	P(C <sub>ox1</sub> )	0.432	C <sub>1</sub>
43	P(C <sub>ox2</sub> )	0.602	C <sub>2</sub>
44	P(N)	0.106	N <sub>3</sub>
45	P(C) <sub>x</sub>	0.545	C <sub>6</sub>
46	P(C) <sub>x, o</sub>	0.432	C <sub>7</sub>
47	P(C) <sub>1</sub>	0.653	C <sub>22</sub>
48	P(C) <sub>1, o</sub>	0.432	C <sub>23</sub>
49	P(C) <sub>1, p</sub>	0.502	C <sub>25</sub>
50	T(C <sub>ox</sub> —C <sub>ox</sub> )	0.085	C <sub>1</sub> —C <sub>2</sub>
51	T(C <sub>ox</sub> =N)	0.491	C <sub>1</sub> —N <sub>4</sub>
52	T(C <sub>ox</sub> —C <sub>x</sub> )	0.695	C <sub>1</sub> —C <sub>6</sub>
53	T(C <sub>ox</sub> —N)	0.037	C <sub>2</sub> —N <sub>3</sub>
54	T(N—C <sub>1</sub> )	0.695	N <sub>3</sub> —C <sub>22</sub>
55	T(N—O)	0.062	N <sub>4</sub> —O <sub>12</sub>
56	T(C—C) <sub>x</sub>	0.278	C <sub>6</sub> —C <sub>7</sub>
57	T(C—C) <sub>1</sub>	0.272	C <sub>22</sub> —C <sub>23</sub>
58	T(C—CH <sub>3</sub> )	0.211	C <sub>25</sub> —C <sub>32</sub>
59	F(C <sub>ox</sub> —C <sub>ox</sub> /C <sub>ox</sub> =N)	0.003	C <sub>1</sub> —C <sub>2</sub> /C <sub>1</sub> —N <sub>4</sub>

No	Force constant		Internal coordinate
60	$F(C_{ox}=N/C_{ox}-C_x)$	0.065	$C_1-N_4/C_1-C_6$
61	$F(C_{ox}=N/C_{ox}=N-O)$	0.055	$C_1-N_4/C_1-N_4-O_{12}$
62	$F(C_{ox}-N/N-C_1)$	-0.175	$C_2-N_3/N_3-C_{22}$
63	$F(C_{ox}-N/C_{ox}-N-C_1)$	1.602	$C_2-N_3/C_2-N_3-C_{22}$
64	$F(C_x-C_x/C_x-C_x)$	1.276*	$C_6-C_7/C_6-C_{11}$
65	$F(C_x-C_x/C_{ox}-C_x-C_x)$	-0.066*	$C_6-C_7/C_1-C_6-C_7$
66	$F(C_x-C_x/C_{ox}-C_x-C_x)$	-0.086	$C_6-C_7/C_1-C_6-C_{11}$
67	$F(C_x-C_x/C_x-C_x-C_x)$	-0.609*	$C_6-C_7/C_7-C_6-C_{11}$
68	$F(C_x-C_x/C_x-C_x-H)$	0.101*	$C_8-C_9/C_9-C_8-H_{18}$
69	$F(C_x-C_x/C-C-Cl)$	0.197	$C_8-C_9/C_8-C_9-Cl_{19}$
70	$F(C_x-C_x/C-Cl)$	0.410	$C_8-C_9/C_9-Cl_{19}$
71	$F(C-Cl/C-C-Cl)$	0.654	$C_9-Cl_{19}/C_8-C_9-Cl_{19}$
72	$F(C-CH_3/C-C-CH_3)$	0.175*	$C_{25}-C_{32}/C_{24}-C_{25}-C_{32}$
73	$F(C-CH_3/C-C_1-H_1)$	0.266*	$C_{25}-C_{32}/C_{25}-C_{32}-H_{33}$
74	$F(C_{ox}/C_x)$	0.083	$C_1/C_6$
75	$F(C_x/C_x)$	-0.065	$C_7/C_8$
76	$F(C_x/C_x-C_x)$	-0.082*	$C_7/C_6-C_7$
77	$F(C_x/C_1-C_1)$	-0.087	$C_{25}/C_{24}-C_{25}$
78	$F(C_x-C_x/C_x-C_x)$	-0.036	$C_6-C_7/C_7-C_8$
79	$F(C_x-C_x/C_x-C_x)$	0.890*	$C_9-C_{10}/C_6-C_{11}$
80	$F(C_x-C_x/C_x-C_x)$	0.279	$C_6-C_7/C_9-C_{10}$
81	$F(C_1-C_1-C_1/C_1-C_1-CH_3)$	-0.064	$C_{23}-C_{24}-C_{25}/C_{24}-C_{25}-C_{32}$
82	$F(C_1/C_1)$	-0.069	$C_{25}/C_{26}$
83	$F(C_1-H_1/C_1-H_1)$	0.073	$C_{32}-H_{33}/C_{32}-H_{34}$
84	$F(C_{ox}-N-H/H-N-C_1)$	0.167*	$C_2-N_3-H_{14}/H_{14}-N_3-C_{22}$
85	$F(C_{ox}-N-H/C_{ox}-N-C_1)$	-0.574*	$C_2-N_3-H_{14}/C_2-N_3-C_{22}$
86	$F(C_{ox}-C_{ox}/C_{ox}-N)$	0.432*	$C_1-C_2/C_2-N_3$
87	$F(C_{ox}-C_{ox}/C_{ox}-C_x)$	-0.051*	$C_1-C_2/C_1-C_6$
88	$F(C_1-C_1/C_1-C_1)$	-0.122*	$C_{24}-C_{25}/C_{25}-C_{26}$

†The force constants are given by K (stretching), H (in-plane bending), P (out-of-plane bending), T (torsion), F (interaction).

‡Bond stretching and bond-bond interaction constants are in  $\text{mdyn}/\text{\AA}$ , angle bending force constants are in  $\text{mdyn } \text{\AA}/\text{rad}^2$ , and bond-angle interaction constants are in  $\text{mdyn}/\text{rad}$ .

§ x: chlorobenzene group, t: methylbenzene group, ox: oxime group, o: ortho, m: meta, p: para

\* Included in the fit.

The frequencies observed in the IR spectra along with their relative intensities, the corresponding calculated frequencies together with the respective potential energy distribution (PEDs) and mode assignments are collected in Table-3. Here, the normal mode description following each fundamental in the last column is due to Wilson<sup>29</sup>. As seen from Table-3, the agreement between experimental and calculated frequencies for pTpCPG is good. The calculated frequencies do not

TABLE-3  
OBSERVED AND CALCULATED PLANAR AND NON-PLANAR FUNDAMENTALS ( $\text{cm}^{-1}$ ) OF pTpCPG\*

Species	Observed Wavenumber (IR)	Calculated Wavenumber (IR-Raman)	Assignments (PED, %)
A'	3367 w	3367	NH stretch [ $\nu(\text{NH})$ ][99K(N—H)]
A'	3307 vw	3307	OH stretch [ $\nu(\text{OH})$ ][100 K(O—H)]
A'	3307 vw	3307	OH stretch [ $\nu(\text{OH})$ ][100 K(O—H)]
A'	3060 w	3066	CH stretch [ $\nu(\text{CH})$ ][98 K(C—H) <sub>x</sub> ]
A'	3060 w	3060	CH stretch [ $\nu(\text{CH})$ ][99 K(C—H) <sub>x</sub> ]
A'	3060 w	3060	CH stretch [ $\nu(\text{CH})$ ][99 K(C—H) <sub>x</sub> ]
A'	3060 w	3060	CH stretch [ $\nu(\text{CH})$ ][99 K(C—H) <sub>x</sub> ]
A'	3037 w	3043	CH stretch [ $\nu(\text{CH})$ ][98 K(C—H) <sub>i</sub> ]
A'	3037 w	3037	CH stretch [ $\nu(\text{CH})$ ][99 K(C—H) <sub>i</sub> ]
A'	3037 w	3037	CH stretch [ $\nu(\text{CH})$ ][99 K(C—H) <sub>i</sub> ]
A'	3037 w	3037	CH stretch [ $\nu(\text{CH})$ ][99 K(C—H) <sub>i</sub> ]
A'	2920 w	2916	CH stretch (methyl), asymmetric [ $\nu(\text{CH})$ ][99 K(C—H)]
A''	2920 w	2915	CH stretch (methyl), asymmetric [ $\nu(\text{CH})$ ][99 K(C—H)]
A'	2860 w	2868	CH stretch (methyl), symmetric [ $\nu(\text{CH})$ ][97 K(C—H)]
A'	1780 w	1777	NH bend [ $\nu(\text{NH})$ ][21 H(H—N—C <sub>1</sub> ) + 20 H(C <sub>ox</sub> —N—H)]
A'	1713 w	1719	CN stretch [ $\nu(\text{C}=\text{N})$ ][53 K(C=N) + 16 K(C—C) <sub>x</sub> ]

Observed Wavenumber (IR)	Calculated Wavenumber (IR-Raman)	Assignments (PED, %)
A' 1684 m	1697	CN + CC stretch [ $\nu(\text{CC})$ , $\nu(\text{CN})$ ] + [39 K(C—C) <sub>t</sub> + 21 H(C—C—H) <sub>t</sub> + 18 K(C=N)]
A' 1675 m	1669	CN + CC stretch [ $\nu(\text{CC})$ , $\nu(\text{CN})$ ][24 K(C—C) <sub>t</sub> + 20 K(C—C) <sub>x</sub> + 14 K(C=N)]
A' 1639 s	1645	CC stretch [ $\nu(\text{CC})$ ] 35 K(C—C) <sub>t</sub> + 33 H(C—C—H) <sub>t</sub> ]
A' 1611 s	1610	CC stretch [ $\nu(\text{CC})$ ] [49 K(C—C) <sub>x</sub> + 17 K(C—C) <sub>t</sub> + 14 H(C—C—H) <sub>x</sub> ]
A' 1592 s	1600	CC stretch [ $\nu(\text{CC})$ ][77 K(C—C) <sub>t</sub> + 17 H(C—C—H) <sub>t</sub> ]
A' 1592 s	1595	CC stretch [ $\nu(\text{CC})$ ][54 K(C—C) <sub>x</sub> + 21 K(C—C) <sub>t</sub> + 21 H(C—C—H) <sub>x</sub> ]
A' 1576 w	1571	CC stretch [ $\nu(\text{CC})$ ][41 K(C—C) <sub>x</sub> + 34 H(C—C—H) <sub>x</sub> ]
A' 1564 w	1564	CC stretch [ $\nu(\text{CC})$ ][58 K(C—C) <sub>x</sub> + 11 K(C—C) <sub>t</sub> + 11 H(C—C—H) <sub>x</sub> ]
A' 1564 w	1557	CC stretch [ $\nu(\text{CC})$ ][51 K(C—C) <sub>t</sub> + 18 K(C—C) <sub>x</sub> + 15 H(C—C—H) <sub>t</sub> ]
A' 1517 s	1517	CC stretch [ $\nu(\text{CC})$ ][29 K(C—C) <sub>t</sub> + 15 K(C—C) <sub>x</sub> + 14 H(C—C—H) <sub>t</sub> ]
A' 1491 s	1478	CC stretch [ $\nu(\text{CC})$ ][46 K(C—C) <sub>x</sub> + 10 K(C—C) <sub>t</sub> ]
A'' 1452 m	1452	CH bend (methyl) [ $\delta(\text{CH})$ ][93 H(H—C <sub>t</sub> —H) + 7 H(C—C <sub>t</sub> —H <sub>t</sub> )]
A' 1429 m	1450	CH bend (methyl), asymmetric [ $\delta(\text{CH})$ ][89 H(H—C <sub>t</sub> —H) + 5 H(C—C <sub>t</sub> —H <sub>t</sub> )]
A' 1393 s	1401	CC stretch [ $\nu(\text{CC})$ ][15 K(C—C) <sub>x</sub> + 15 K(C—C) <sub>t</sub> + 14 K(C <sub>ox</sub> —C <sub>x</sub> ) + 5 K(C <sub>ox</sub> —N) + 6 H(C—C—H) <sub>x</sub> + 6 H(H—N—C <sub>t</sub> )]
A' 1393 s	1400	CC stretch [ $\nu(\text{CC})$ ][50 K(C—C) <sub>x</sub> + 14 K(C <sub>ox</sub> —C <sub>x</sub> )]
A' 1393 s	1395	CH bend (methyl), symmetric [ $\delta(\text{CH})$ ][52 H(C—C <sub>t</sub> —H <sub>t</sub> ) + 43 H(H—C <sub>t</sub> —H)]



Species	Observed Wavenumber (IR)	Calculated Wavenumber (IR-Raman)	Assignments (PED, %)
A'	1326 m	1343	CC stretch [ $\nu(\text{CC})$ ][36 K(C—C) <sub>t</sub> + 12 F(C <sub>x</sub> —C <sub>x</sub> /C <sub>x</sub> —C <sub>x</sub> ) + 10 H(C—C—H) <sub>d</sub> ]
A'	1310 m	1306	CH i.p. bend [ $\delta(\text{CH})$ ][25 H(C—C—H) <sub>t</sub> + 23 H(C—C—H) <sub>x</sub> + 18 K(C—C) <sub>x</sub> ]
A'	1297 w	1302	CH i.p. bend [ $\delta(\text{CH})$ ][47 H(C—C—H) <sub>t</sub> + 25 H(C—C—H) <sub>x</sub> ]
A'	1282 m	1278	CH i.p. bend [ $\delta(\text{CH})$ ][34 H(C—C—H) <sub>x</sub> + 17 H(N—O—H) + 18 K(C—C) <sub>x</sub> ]
A'	1260 w	1254	CH i.p. bend [ $\delta(\text{CH})$ ][53 H(C—C—H) <sub>t</sub> + 13 K(C—CH <sub>3</sub> ) + 13 K(N—C <sub>l</sub> ) + 11 H(N—O—H)]
A'	1241 w	1228	OH i.p. bend [ $\delta(\text{OH})$ ][78 H(N—O—H) + 9 K(N—O)]
A'	1210 w	1204	OH i.p. bend [ $\delta(\text{OH})$ ][39 H(N—O—H) + 11 K(C—C) <sub>x</sub> ]
A''	1210 w	1209	CH o.p. bend [ $\gamma(\text{C})$ ][63 P(C) <sub>t</sub> + 12 T(C—C) <sub>t</sub> + 12 F(C <sub>x</sub> /C <sub>x</sub> )]
A''	1178 w	1163	CH o.p. bend [ $\gamma(\text{C})$ ][66 P(C) <sub>x</sub> + 18 T(C—C) <sub>x</sub> ]
A'	1130 w	1128	CH i.p. bend [ $\delta(\text{CH})$ ][54 H(C—C—H) <sub>x</sub> + 14 K(C—C) <sub>x</sub> ]
A'	1130 w	1124	CH i.p. bend [ $\delta(\text{CH})$ ][64 H(C—C—H) <sub>t</sub> + 22 K(C—C) <sub>t</sub> ]
A'	1112 w	1118	CH i.p. bend [ $\delta(\text{CH})$ ][47 H(C—C—H) <sub>x</sub> + 15 K(C—O) <sub>x</sub> ]
A''	1112 w	1113	CH o.p. bend [ $\gamma(\text{C})$ ][69 P(C) <sub>x</sub> + 24 T(C—C) <sub>x</sub> ]
A'	1093 s	1109	CH i.p. bend [ $\delta(\text{CH})$ ][62 H(C—C—H) <sub>x</sub> + 25 K(C—O) <sub>x</sub> ]
A'	1093 s	1108	CH i.p. bend [ $\delta(\text{CH})$ ][62 H(C—C—H) <sub>t</sub> + 30 K(C—C) <sub>t</sub> ]
A''	1075 w	1069	CH bend (methyl) [ $\delta(\text{CH})$ ][39 H(C—C—H) <sub>d</sub> + 37 P(C) <sub>t</sub> ]
A'	1042 w	1028	CH i.p. bend [ $\delta(\text{CH})$ ][29 H(C—C—H) <sub>x</sub> + 39 H(C—C—C) <sub>x</sub> + 19 K(C—C) <sub>x</sub> ]
A'	1020 w	1028	CH i.p. bend [ $\delta(\text{CH})$ ][34 H(C—C—H) <sub>t</sub> + 29 H(C—C—C) <sub>t</sub> + 26 K(C—C) <sub>t</sub> ]

Species	Observed Wavenumber (IR)	Calculated Wavenumber (IR-Raman)	Assignments (PED, %)
A'	1013 m	1019	NO stretch [ $\nu(\text{NO})$ ][55 K(N—O) + 16 K(C <sub>ox</sub> —C <sub>ox</sub> )]
A''	994 m	1004	CH o.p. bend. [ $\gamma(\text{C})$ ][72 P(C) <sub>t</sub> + 15 T(C—C) <sub>l</sub> ]
A'	972 m	972	CH bend (methyl) [ $\delta(\text{CH})$ ][83 H(C—C <sub>t</sub> —H <sub>l</sub> )]
A''	957 s	955	CH o.p. bend. [ $\gamma(\text{C})$ ][51 P(C) <sub>t</sub> + 40 H(C—C <sub>t</sub> —H <sub>l</sub> ) + 14 T(C—C) <sub>l</sub> ]
A'	904 m	910	NO stretch [ $\nu(\text{NO})$ ][36 K(N—O) + 12 K(C—C) <sub>l</sub> ]
A''	890 m	890	CH o.p. bend. [ $\gamma(\text{C})$ ][81 P(C) <sub>x</sub> + 10 T(C—C) <sub>x</sub> ]
A'	851 w	850	CN stretch [ $\nu(\text{C}_{\text{ox}}\text{N})$ ][48 K(C <sub>ox</sub> —N) + 32 F(C <sub>ox</sub> —N/C <sub>ox</sub> —N—C <sub>l</sub> ) + 15 H(C <sub>ox</sub> —N—C <sub>l</sub> )]
A'	830 s	824	NO stretch [ $\nu(\text{NO})$ ][23 K(N—O) + 22 K(C—CH <sub>3</sub> ) + 36 K(C—C) <sub>l</sub> ]
A''	818 s	814	CH o.p. bend. [ $\gamma(\text{C})$ ][87 P(C) <sub>x</sub> + 10 T(C—C) <sub>x</sub> ]
A''	799 m	795	CN torsion [ $\tau(\text{CN})$ ][49 T(N—C <sub>l</sub> ) + 30 P(C) <sub>t</sub> + 18 P(N)]
A''	761 m	783	CC o.p. bend. [ $\gamma(\text{C})$ ][29 P(C <sub>ox</sub> 2) + 22P(C <sub>ox</sub> 1) + 26 P(C) <sub>x</sub> ]
A''	731 m	739	CH o.p. bend. [ $\gamma(\text{C})$ ][80 P(C) <sub>t</sub> + 18 T(C—C) <sub>l</sub> ]
A'	721 m	734	CCl stretch [ $\nu(\text{CCl})$ ][25 K(C—Cl) + 36 K(C—C) <sub>x</sub> + 16 H(C—C—C) <sub>x</sub> ]
A'	704 w	720	NO stretch [ $\nu(\text{NO})$ ][17 K(N—O) + 12 K(C—CH <sub>3</sub> ) + 13 K(N—C <sub>l</sub> ) + 13 H(C—C—C) <sub>l</sub> ]
A''	681 m	692	CC o.p. bend. [ $\gamma(\text{C})$ ][26 P(C <sub>ox</sub> 2) + 22 P(C) <sub>x</sub> + 12 T(C <sub>ox</sub> =N) + 12 T(C—C) <sub>x</sub> ]
A''	661 w	663	CH o.p. bend. [ $\gamma(\text{C})$ ][40 P(C) <sub>t</sub> + 40 T(C—C) <sub>l</sub> ]
A'	630 w	621	CCl stretch [ $\nu(\text{CCl})$ ][24 K(C—Cl) + 13 H(C <sub>ox</sub> =N—O) + 13 H(C—C—C) <sub>x</sub> ]
A''	576 m	589	CH o.p. bend. [ $\gamma(\text{C})$ ][28 P(C) <sub>x</sub> + 25 T(C—C) <sub>x</sub> + 18 T(C <sub>ox</sub> =N) + 2014 P(C <sub>ox</sub> 1)]

